KEVOFLOX-BH



LEVOFLOXACIN HEMIHYDRATE,
TIAMULIN HYDROGEN FUMARATE &
BROMHEXINE HYDROCHLORIDE
ORAL SOLUTION

Composition:

Levofloxacin Hemihydrate I.P Tiamulin Hydrogen Fumarate USP Bromhexine Hydrochloride IP Aqueous Base 10% w/v 1.0% w/v 1.5% w/v q.s.



Indications

- Bacterial infections like infectious Coryza, Fowl Choera, Necrotic Enteritis, Colibacillosis, Salmonellosis, and other mixed non-specific bacterial infections.
- Sever respiratory infections like chronic Respiratory Disease (CRD) & CCRD.
- ECM associated with yolk sac infection, MDR (Multi Drug Resistance) cases.
- Secondary bacterial complication after viral infections.

Storage:

Store in a cool dry and dark place. In case of crystallization keep bottle in warm water for 30 minutes

Administration & Dosage:

Mix 1 ml. for 10 kgs of live body weight
To be administered continuously for 3-5 days
once Daily in drinking water depending upon
the severity of infection or as directed by the veterinarian

Withdrawal Period:

Meat: 28 Days, Eggs: 7 days At a dose of 1 ml per 10 kg body weight, 1 mg Tiamulin is made available per kg body weight, which is well within safety limit for use with any coccidiostat.

